

# THE HURWITZ EQUIVALENCE PROBLEM IS UNDECIDABLE

E. LIBERMAN<sup>1</sup> AND M. TEICHER

ABSTRACT. In this paper, we prove that the Hurwitz equivalence problem for 1-factorizations in  $F_2 \oplus F_2$  is undecidable, and as a consequence, the Hurwitz equivalence problem for  $\Delta^2$ -factorizations in the braid groups  $B_n, n \geq 5$  is also undecidable.

## INTRODUCTION

It has long been conjectured (e.g [12]) that the Hurwitz equivalence problem is undecidable. In this paper we present a proof to this conjecture.

This problem relates to Algebraic Geometry as follows: There is a well defined construction (see [12]) which attaches to any projective curve a  $\Delta^2$ -factorization of the braid group, called the *braid monodromy factorization*. This gives rise to the definition of the braid monodromy type (BMT) of projective curves: Two curves have the same BMT if their braid monodromy factorizations are Hurwitz equivalent up to at most one global conjugation. In the same paper it is shown that two cuspidal curves are isotopic if and only if their BMT are equal. This profound theorem invites the following question: Does there exist a finite algorithm which recognizes whether two braid monodromy factorizations belong to the same braid monodromy type? In order to answer this question, we first ask a slightly simpler question: Does there exist a finite algorithm which determines whether two  $\Delta^2$ -factorizations are Hurwitz-equivalent? In this paper we intend to show that the answer

---

*Date:* February 2, 2008.

<sup>1</sup>Partially supported by the Emmy Noether Research Institute for Mathematics, Bar-Ilan University, and the Minerva Foundation, Germany, and by the Excellency Center “Group Theoretic Methods in the study of Algebraic Varieties” of the National Science Foundation of Israel.

to this question is negative. This problem, of determining whether two factorizations are Hurwitz-equivalent, has been discussed in many papers, but only partial results have been achieved.

In this paper we shall find a connection between the Hurwitz-equivalence problem and the word problem of finitely presented groups. The word problem is very well known, and in ([5],[11]) it is shown that there exist finitely presented groups whose word problem is unsolvable. In ([2],[1],[13]) it is shown that determining whether the word problem of a given group is solvable is itself undecidable. In [6] it is shown further, that determining whether a group's word problem is solvable is unrecognizable, and there is no uniform partial algorithm which solves the word problem for all the finitely presented groups whose word problem is solvable. In [8] Higman proves the existence of a universal finitely presented group  $K$ , such that there exists a Turing machine which, given a finitely presented group, finds a finite subset of  $K$  which generates a subgroup isomorphic to the given group. For more information on the word problem and decision problems in group theory, see for example [10].

This paper is organized as follows: In chapter 1 we shall present some well known definitions and results which we intend to investigate. In chapter 2 we shall study the structure of the Hurwitz stabilizer, and the effect of the Hurwitz braid action on direct products. In chapter 3 we shall define a new structure and study the connection between the Hurwitz equivalence problem and the problem of finding a solution to the equation  $H^Y = H_1$ , where  $Y$  comes from a specified normal subgroup. In chapter 4 we present our main theorem, which connects between the word problem of an arbitrary group and the 1-factorizations of the braid group. Finally, in chapter 5, we shall prove that the Hurwitz-equivalence problem is undecidable for  $F_2 \oplus F_2$ , and as an important consequence, that the Hurwitz-equivalence problem for  $\Delta^2$ -factorizations in the braid group  $B_n, n \geq 5$  is undecidable.

## 1. PRELIMINARIES

The braid group is the group

$$\left\langle \sigma_1, \dots, \sigma_{n-1} \left| \begin{array}{ll} \sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1} \\ \sigma_i \sigma_j = \sigma_j \sigma_i & \text{if } |i - j| \geq 2 \end{array} \right. \right\rangle$$

There is a natural homomorphism  $\pi : B_n \rightarrow S_n$ , defined by mapping  $\sigma_i$  to the transposition  $(i, i + 1)$ . The permutation  $\pi(b)$  of a braid  $b$  is called the braid permutation. There is a normal subgroup  $PB_n$  of  $B_n$ , consisting of those braids whose braid permutation is trivial. It is shown in [7] that the generators of  $PB_n$  are

$$A_{ij} = \sigma_i^{-1} \cdots \sigma_{j-2}^{-1} (\sigma_{j-1}^2) \sigma_{j-2} \cdots \sigma_i$$

, with  $1 \leq i < j \leq n$ .

**Theorem 1.1.** [3] *Let  $b$  be a braid in  $B_n$ , and  $\pi$  any braid with the same braid permutation. Then  $b$  can be written as  $\pi U_n \dots U_1$ , where each  $U_i$  is generated only by generators of the form  $A_{ik}$ .*

More information on the pure braid groups and braid combing – the process of rewriting a braid in this normal form – can be found in [3] and [4].

In this paper we shall be particularly interested in a (right) group action called the *Hurwitz braid action*.

**Definition 1.2.** *Let  $G$  be a group. We call a vector  $F = (f_1, \dots, f_n) \in G^n$  a  $g$ -factorization of length  $n$  if its product  $m(F) := \prod_{i=1}^n f_i$  is equal to  $g$ .*

*Let  $n$  be a natural number, and  $g \in G$ . The Hurwitz braid action is the (right) group action of  $B_n$  on the set of  $g$ -factorizations of length  $n$ , defined as follows: Let  $F = (f_1, \dots, f_n)$  be a factorization then*

$$\begin{aligned} (f_1, \dots, f_n) \sigma_i &= (f_1, \dots, f_{i-1}, f_{i+1}, f_{i+1}^{-1} f_i f_{i+1}, f_{i+2}, \dots, f_n) \\ (f_1, \dots, f_n) \sigma_i^{-1} &= (f_1, \dots, f_{i-1}, f_i f_{i+1} f_i^{-1}, f_i, f_{i+2}, \dots, f_n) \end{aligned}$$

From now on, when we refer to a group action, we mean the Hurwitz braid action.

**Definition 1.3** (Hurwitz equivalence). *Two factorizations are said to be Hurwitz equivalent if they belong to the same group orbit. We shall denote this equivalence relation by  $\cong_H$ .*

Note that nothing crucial would change if we chose to define the equivalence through a left group action.

**Definition 1.4.** *Let  $F = (f_1, \dots, f_n)$  be a factorization of length  $n$ , and  $K = (k_1, \dots, k_m)$  a factorization of length  $m$ . We denote by*

$F \otimes K$  the concatenation of these two factorizations, i.e.  $F \otimes K = (f_1, \dots, f_n, k_1, \dots, k_m)$ . Evidently this is a factorization of length  $m+n$ .

**Definition 1.5.** Let  $X = (g_1, \dots, g_s)$  and  $Y$  be factorizations. Then  $X^Y := (g_1^{m(Y)}, \dots, g_s^{m(Y)})$  and  $X^{(Y^{-1})} := (g_1^{m(Y)^{-1}}, \dots, g_s^{m(Y)^{-1}})$

We shall now give an example of how the generators  $A_{ij}$  act on a factorization.

**Example 1.6.** Let  $F$  be a factorization. Let  $X$  be the first  $i-1$  elements of  $F$ ,  $a$  be the  $i$ -th element,  $Y$  be the elements between the  $i+1$ -th and the  $j-1$ -th elements,  $b$  be the  $j$ -th element and  $Z$  be the remaining elements of  $F$ , from the  $j+1$ -th element and onward. Then

$$\begin{aligned} (X \otimes a \otimes Y \otimes b \otimes Z)A_{ij} &= (X \otimes a \otimes Y \otimes b \otimes Z)\sigma_i^{-1} \cdots \sigma_{j-2}^{-1}(\sigma_{j-1}^2)\sigma_{j-2} \cdots \sigma_i = \\ (X \otimes Y^{a^{-1}} \otimes a \otimes b \otimes Z)(\sigma_{j-1}^2)\sigma_{j-2} \cdots \sigma_i &= (X \otimes Y^{a^{-1}} \otimes b \otimes a^b \otimes Z)(\sigma_{j-1})\sigma_{j-2} \cdots \sigma_i \\ &= (X \otimes Y^{a^{-1}} \otimes a^b \otimes b^{a^b} \otimes Z)\sigma_{j-2} \cdots \sigma_i = \\ (X \otimes a^b \otimes Y^{a^{-1}a^b} \otimes b^{a^b} \otimes Z) \end{aligned}$$

$$\begin{aligned} (X \otimes a \otimes Y \otimes b \otimes Z)A_{ij}^{-1} &= (X \otimes a \otimes Y \otimes b \otimes Z)\sigma_i^{-1} \cdots \sigma_{j-2}^{-1}(\sigma_{j-1}^{-2})\sigma_{j-2} \cdots \sigma_i = \\ (X \otimes Y^{a^{-1}} \otimes a \otimes b \otimes Z)(\sigma_{j-1}^{-2})\sigma_{j-2} \cdots \sigma_i &= (X \otimes Y^{a^{-1}} \otimes b^{a^{-1}} \otimes a \otimes Z)(\sigma_{j-1})\sigma_{j-2} \cdots \sigma_i \\ &= (X \otimes Y^{a^{-1}} \otimes a^{(b^{a^{-1}})^{-1}} \otimes b^{a^{-1}} \otimes Z)\sigma_{j-2} \cdots \sigma_i = \\ (X \otimes a^{(b^{a^{-1}})^{-1}} \otimes Y^{a^{-1}a^{(b^{a^{-1}})^{-1}}} \otimes b^{a^{-1}} \otimes Z) \end{aligned}$$

**Corollary 1.7.** We can see from this example that  $A_{ij}$  and  $A_{ij}^{-1}$  replace the  $i$ -th element of the factorization with a conjugate of itself, while the other elements are either unchanged or conjugated by an element from the normal closure of the  $i$ -th element in the subgroup generated by the factorization elements.

### 1.1. Recursiveness and Recursive Enumerability.

An *alphabet* is a finite set  $\Sigma$ . Elements of  $\Sigma$  are called *symbols*. *Words* from  $\Sigma$  are finite sequences of symbols from  $\Sigma$ .  $\Sigma^*$  is the set of all words from  $\Sigma$ .

Let  $\Sigma$  be an alphabet, and  $X$  a subset of  $\Sigma^*$ .  $X$  is called a recursively enumerable set if there exists a Turing machine  $T$  such that  $T$

halts on any word from  $X$ , and does not halt otherwise.  $X$  is called a recursive set if there is a Turing machine which outputs 1 on any input from  $X$ , and outputs 0 for inputs from  $\Sigma^* \setminus X$ . Similarly, a decision problem is called solvable if the set of elements whose answer is ‘yes’ is recursive, and recognizable if the set of elements whose answer is ‘yes’ is recursively enumerable.

## 2. PROPERTIES OF THE HURWITZ BRAID ACTION

The following proposition (Proposition 2.1) describes the basic properties of the Hurwitz braid action and the Hurwitz equivalence. The properties are all natural, and are detailed here for the readers convenience.

### Proposition 2.1.

- (1) *Let  $G = (g_1, \dots, g_n)$  and  $K = (k_1, \dots, k_n)$  be factorizations. If  $G \cong_H K$ , then the group generated by the elements of  $G$  is equal to the group generated by the elements of  $K$ , i.e.,  $\langle g_1, \dots, g_n \rangle = \langle k_1, \dots, k_n \rangle$ .*
- (2) *Let  $K$  be a subgroup of  $G$ , and let  $(g_1, \dots, g_n)$  and  $(h_1, \dots, h_n)$  be two factorizations such that  $g_i$  and  $h_i$  are elements from  $K$ . Then  $(g_1, \dots, g_n) \cong_H (h_1, \dots, h_n)$  as factorizations in  $K$  if and only if  $(g_1, \dots, g_n) \cong_H (h_1, \dots, h_n)$  as factorizations in  $G$ .*
- (3) *Let  $(g_1, \dots, g_n)$  be a factorization and  $G = \langle g_1, \dots, g_n \rangle$ . Let  $b$  be a braid with braid permutation  $\pi$ . Assume that  $(g_1, \dots, g_n)b = (h_1, \dots, h_n)$ . Then, for any  $i$ ,  $g_i$  and  $h_{\pi(i)}$  are conjugate in  $G$ .*
- (4) *Let  $h : G_1 \rightarrow G_2$  be a group homomorphism, and let  $F = (f_1, \dots, f_n)$  be a factorization in the group  $G_1$ . Denote  $h(F) = (h(f_1), \dots, h(f_n))$ . Then, for any braid  $b \in B_n$ ,  $[h(F)b] = h[(F)b]$ .*

*Proof.* (1) We first prove the proposition for  $\sigma_i$ . Let  $F = (f_1, \dots, f_n)$  be a factorization, and  $G_1 = \langle f_1, \dots, f_m \rangle$  be the group generated by its elements. Similarly, Let  $G_2$  be the group generated by the elements of  $F\sigma_i = (f_1 \dots f_{i-1}, f_{i+1}, f_{i+1}^{-1}f_i f_{i+1}, f_{i+2}, \dots, f_n)$ , i.e.  $G_2 = \langle f_1 \dots f_{i-1}, f_{i+1}, f_{i+1}^{-1}f_i f_{i+1}, f_{i+2}, \dots, f_n \rangle$ . All the elements besides the  $i$ -th and  $i+1$ -th are equal, so in order to prove equality, it suffices to show that  $f_i, f_{i+1} \in G_2$  and

that  $f_{i+1}, f_{i+1}^{-1}f_i f_{i+1} \in G_1$ . Evidently  $f_{i+1}$  belongs to both groups. Now, since  $f_i$  and  $f_{i+1}$  are in  $G_1$ , so is  $f_{i+1}^{-1}f_i f_{i+1}$ . Similarly, since  $f_{i+1}$  and  $f_{i+1}^{-1}f_i f_{i+1}$  belong to  $G_2$ , so does  $f_i = (f_{i+1})(f_{i+1}^{-1}f_i f_{i+1})(f_{i+1}^{-1})$ . This completes the proof on the generators of the braid group, and the proof can easily be completed by induction.

- (2) From (1) we see that if  $F = (f_1, \dots, f_n)$  is a factorization such that  $f_i \in K$ , then all the elements in the factorizations belonging to the orbit of  $F$  also belong to  $K$ . Now by hypothesis  $K$  is a subgroup of  $G$ , so the action of  $\sigma_i$  on  $F$  is the same, whether we consider  $F$  a factorization in  $G$  or in  $K$ . Therefore, regardless of our point of view, the group orbit of  $F$  remains the same. Since by definition  $F \cong_H F_1$  if  $F_1$  is in the group orbit as  $F$ , we have completed the proof.
- (3) We first give a proof for the generators  $\sigma_i$ . All the elements in the factorization other than the  $i$ -th and  $(i+1)$ -th elements are left unchanged by the braid action, and the braid permutation  $(i, i+1)$  also leaves them fixed, so the theorem holds trivially. It remains to check the  $i$ -th and  $(i+1)$ -th elements.  $\sigma_i$  maps  $(f_i, f_{i+1})$  to  $(f_{i+1}, f_{i+1}^{-1}f_i f_{i+1})$ , and we see clearly that the  $(i+1)$ -th element of the first factorization is equal (and hence conjugate) to the  $i$ -th element in the second one, and that the  $i$ -th element of the first factorization is conjugate to the  $(i+1)$ -th element of the second one. The proof can now be completed by a straightforward induction, bearing in mind that from (1) we know that every factorization that is Hurwitz equivalent to our factorization has its elements in  $G$  too.
- (4) Again, we shall start by proving the proposition for  $\sigma_i$ . We know that  $\sigma_i$  affects only the  $i$ -th and  $(i+1)$ -th elements of the factorization, so we need only consider them. We know that the  $i$ -th and  $(i+1)$ -th elements of  $h(F)$  are  $h(f_i)$  and  $h(f_{i+1})$  respectively, so  $(h(f_i), h(f_{i+1}))\sigma_i = (h(f_{i+1}), h(f_{i+1}^{-1}f_i f_{i+1}))$ . since  $h$  is a group homomorphism, this is equal to  $(h(f_{i+1}), h(f_{i+1}^{-1}f_i f_{i+1}))$ , which are precisely the  $i$ -th and  $(i+1)$ -th elements of  $h[(F)\sigma_i]$ . Once again, the proof is easily completed by induction.

□

**Lemma 2.2.** *Let  $h : G_1 \rightarrow G_2$  be a homomorphism, and  $F$  a factorization of  $G_1$ . Then  $\text{stab}(F) \subseteq \text{stab}(h(F))$ .*

*Proof.* Let  $b \in \text{stab}(F)$ , which means that  $Fb = F$ . From Lemma 2.1 we have that  $[h(F)]b = h[(F)b]$ . Putting both together we obtain  $[h(F)]b = h(F)$ , which implies that  $b \in \text{stab}(h(F))$ .  $\square$

**Lemma 2.3.** *Let  $V = (x_1, \dots, x_r)$  be a factorization such that  $x_i \neq 1$  for all  $i$ , and let  $S$  be the stabilizer of  $V$ . Let  $\varphi : B_r \rightarrow B_{n+r}$  be defined by  $\sigma_i \mapsto \sigma_{i+n}$ . Let*

$$V_1 = (\underbrace{1, \dots, 1}_n, x_1, \dots, x_r)$$

*Then, the stabilizer of  $V_1$  is  $S_1 = \langle \{\sigma_i \mid i < n\}, \{A_{jk} \mid j < n\}, \varphi(S) \rangle$ .*

*Furthermore, we can write any element of the stabilizer in the form  $CBA$ , where  $A \in \langle \sigma_i \mid i < n \rangle$ ,  $B \in \langle A_{jk} \mid j < n \rangle$  and  $C \in \varphi(S)$ .*

*Proof.*  $S_1 \subseteq \text{stab}(V_1)$ .

We shall show this by considering the three types of braids in  $S_1$ .

- $\sigma_i, i < n$ .  $\sigma_i$  affects only the  $i$ -th and  $(i+1)$ -th elements, which in our case are  $(1, 1)$ . The operation of  $\sigma_i$  on these gives  $(1, 1^{-1} \cdot 1 \cdot 1) = (1, 1)$ .
- $A_{jk}, j < n$ . From example 1.6 we know that if  $V$  is a factorization with the same notation as in the example, then  $(V)A_{jk} = (X \otimes a^{ba^{-1}} \otimes Y^{a^{-1}a^{ba^{-1}}} \otimes b^{a^{-1}} \otimes Z)$ . However, in our case  $a = 1$ , so we get

$$(X \otimes 1^{b*1^{-1}} \otimes Y^{1^{-1}1^{b*1^{-1}}} \otimes b^{1^{-1}} \otimes Z) = (X \otimes 1 \otimes Y \otimes b \otimes Z) = V$$

- $\varphi(S)$ . The elements of  $\varphi(S)$  act only on the last  $r$  elements of the factorizations, and do so the same way  $S$  acts on  $V$ . Therefore,  $\varphi(S) \subseteq \text{stab}(V_1)$ .

$$\text{stab}(V_1) \subseteq S_1.$$

Let  $b$  is a braid in the stabilizer. From Theorem 1.1 we know that we can write  $b$  as  $b = \pi F_{r+n-1} \cdots F_1$ , where  $\pi$  is an arbitrary braid with the same braid permutation as  $b$ , and the  $F_i$  is generated by braids of the form  $A_{ik}$ , where  $i$  is fixed and  $i < k$ . By hypothesis we know that  $x_i \neq 1$  for all  $i$ , so the  $x_i$ s cannot be conjugate to 1. Therefore,  $b$ 's braid permutation must permute the elements  $1, \dots, n$  among themselves and

the elements  $n+1, \dots, n+r$  among themselves. This means that  $\pi$  can be factored as  $\pi_1\pi_2$ , where  $\pi_1$  is generated by braids from  $\{\sigma_i \mid i > n\}$  and  $\pi_2$  is generated by braids from  $\{\sigma_i \mid i < n\}$ .

Now, since the pure braid group is a normal subgroup and  $F_{r+n-1} \cdots F_1 \in PB_n$ , then  $\pi_2 F_{r+n-1} \cdots F_1 = P\pi_2$ , for some  $P \in PB_n$ . This means that  $P$  can be written as  $F'_{r+n-1} \cdots F'_1$ , so we have  $\pi_2 F_{r+n-1} \cdots F_1 = F'_{r+n-1} \cdots F'_1 \pi_2$ . Therefore,  $b = \pi_1 \pi_2 F_{r+n-1} \cdots F_1 = \pi_1 (F'_{r+n-1} \cdots F'_1) \pi_2$ .

We know that  $b$  is in the stabilizer, so  $(V_1)b = V_1$ , which means  $(V_1)\pi_1(F'_{r+n-1} \cdots F'_1)\pi_2 = V_1$ , or in other words  $(V_1)(\pi_1 F'_{r+n-1} \cdots F'_n)(F'_{n-1} \cdots F'_1 \pi_2) = V_1$ . The braid permutation of  $(\pi_1 F'_{r+n-1} \cdots F'_n)$  acts only on the first  $n$  elements, hence the first  $n$  elements are conjugate to 1, which means they are equal to 1. The last  $r$  elements are conjugated among themselves and therefore they are not equal to 1. Therefore, we can implement the part of the lemma which we proved above on  $(V_1)(\pi_1 F'_{r+n-1} \cdots F'_n)$ , and conclude that  $F'_{n-1} \cdots F'_1 \pi_2$  is in the stabilizer of  $(V_1)(\pi_1 F'_{r+n-1} \cdots F'_n)$ .

This in turn implies that  $(V_1)(\pi_1 F'_{r+n-1} \cdots F'_n) = V_1$ , so therefore  $(\pi_1 F'_{r+n-1} \cdots F'_n)$  is in the stabilizer of  $V_1$ . Note also that it is generated by braids of the form  $\{\sigma_i \mid i > n\}$ , and therefore it is in  $\varphi(S)$ . In summary, we have shown that the braid  $b$  can be written as  $[\pi_1 F'_{r+n-1} \cdots F'_n][F'_{n-1} \cdots F'_1]\pi_2$ , where  $\pi_2 \in \langle \{\sigma_i \mid i < n\} \rangle$ ,  $[F'_{n-1} \cdots F'_1] \in \langle \{A_{jk} \mid j < n\} \rangle$ ,  $[\pi_1 F'_{r+n-1} \cdots F'_n] \in \varphi(S)$ , which completes the proof.  $\square$

**Definition 2.4.** Let  $G = A \oplus B = \{(a, b) \mid a \in A, b \in B\}$ . Let  $V = (v_1, \dots, v_n)$  be a factorization in  $A$  and  $W = (w_1, \dots, w_2)$  a factorization in  $B$ . We denote by  $V \oplus W$  the factorization of  $A \oplus B$  whose  $i$ -th element is  $(v_i, w_i)$ . In other words,  $V \oplus W = ((v_1, w_1), \dots, (v_n, w_n))$ .

**Lemma 2.5.** Let  $b$  be a braid, then  $(V \oplus W)b = (V)b \oplus (W)b$ .

*Proof.* First we shall prove the lemma for  $\sigma_i$ . We know that  $\sigma_i$  affects only the  $i$ -th and  $(i+1)$ -th elements of the factorization. Let us write the  $i$ -th element of  $V \oplus W$  as  $(a_1, b_1)$  and the  $(i+1)$ -th element as  $(a_2, b_2)$ . Now,  
 $((a_1, b_1), (a_2, b_2))\sigma_1 = ((a_2, b_2), (a_2, b_2)^{-1}(a_1, b_1)(a_2, b_2))$



$= ((a_2, b_2), (a_2^{-1}a_1a_2, b_2^{-1}b_1b_2))$ , which are precisely the  $i$ -th and  $(i+1)$ -th elements of  $(V)\sigma_i \oplus (W)\sigma_i$ . A similar argument holds for  $\sigma_i^{-1}$ . The proof for arbitrary braids follows by induction.  $\square$

**Lemma 2.6.**  *$V \oplus W \cong_H V_1 \oplus W$  if and only if there exists a braid  $b \in \text{stab}(W)$  such that  $(V)b = V_1$ .*

*Proof.* By definition,  $V \oplus W \cong_H V_1 \oplus W$  if and only if there is a braid  $b$  such that  $(V \oplus W)b = V_1 \oplus W$ . However, from lemma 2.5 we know that  $(V \oplus W)b = (V)b \oplus (W)b$ , so our requirement becomes  $(V)b \oplus (W)b = V_1 \oplus W$ . This implies that  $(W)b = W$ , i.e. that  $b \in \text{stab}(W)$ , and that  $(V)b = V_1$ .  $\square$

### 3. ON THE SET OF FACTORIZATIONS

In this section we construct a new structure on the set of factorizations and study its properties.

**Definition 3.1.** *Let  $G = G_1 \oplus G_2$ . Let  $R_1, \dots, R_n, W_1, \dots, W_m, H \in G_1$  and  $X_1, \dots, X_{m+2} \in G_2$ . Let  $R$  be the factorization  $(R_1, \dots, R_n)$ ,  $W$  the factorization  $(W_1, \dots, W_m)$  and  $X$  the factorization  $(X_1, \dots, X_{m+2})$ . We define  $P_X(R, W, H)$  to be the following factorization on  $G$ :*

$$P_X(R, W, H) = ((R_1, 1), \dots, (R_n, 1), (W_1, X_1), \dots, (W_m, X_m), (H^{-1}, X_{m+1}), (H, X_{m+2}))$$

We will now present two theorems which give necessary and sufficient conditions for the equivalence of the Hurwitz equivalence problem and the problem of finding an element  $Y$  inside a normal subgroup such that  $H^Y = H_1$ .

**Theorem 3.2.** *Let  $R$  be the factorization  $(R_1, \dots, R_n)$ ,  $W$  be the factorization  $(W_1, \dots, W_m)$  and  $X$  be the factorization  $(X_1, \dots, X_{m+2})$ . Let  $H$  and  $H_1$  be elements of  $G_1$ .*

*Suppose that  $R_1 \cdots R_n = 1$ . Let  $N$  be the normal closure of  $\{R_1, \dots, R_n\}$  inside  $\langle R_1, \dots, R_n, W_1, \dots, W_m \rangle$ , and assume that  $N$  contains a word  $Y$  such that  $H^Y = H_1$ . Then  $P_X(R, W, H) \cong_H P_X(R, W, H_1)$ .*

In order to prove Theorem 3.2, we first prove two lemmas.

**Lemma 3.3.** *Let  $R$  be the factorization  $(R_1, \dots, R_n)$ ,  $W$  be the factorization  $(W_1, \dots, W_m)$  and  $X$  be the factorization  $(X_1, \dots, X_{m+2})$ . Let*

$H$  be an element from  $G_1$ . Suppose also that  $R_1 \cdots R_n = 1$ . Then for any  $j$ :  $P_X(R, W, H) \cong_H P_X(R^{W_j}, W, H)$

*Proof.* We know from Lemma 2.6 that  $P_X(R, W, H) \cong_H P_X(R^{W_j}, W, H)$  if and only if there exists a braid  $b \in B_{m+2}$  in the stabilizer of  $(\underbrace{1, \dots, 1}_n, X_1, \dots, X_{m+2})$ ,

such that  $(R_1, \dots, R_n, W_1, \dots, W_m, H^{-1}, H)b$  is equal to  $(R_1^{W_j}, \dots, R_n^{W_j}, W_1, W_m, H^{-1}, H)$ .

With this in mind, define the braid

$$b_j = (\sigma_{j-1} \cdots \sigma_n)(\sigma_{n-1} \cdots \sigma_1)(\sigma_1 \cdots \sigma_{n-1})(\sigma_{j-1} \cdots \sigma_n)^{-1} \text{ for } j > n.$$

We begin by showing that  $b_j$  is in the stabilizer of  $(\underbrace{1, \dots, 1}_n, X_1, \dots, X_{m+2})$ .

Let  $I = (\underbrace{1, \dots, 1}_n)$ ,  $Y = (X_1, \dots, X_j)$  and  $Z = (X_{j+1}, \dots, X_{m+2})$  and so:

$$\begin{aligned} (\underbrace{1, \dots, 1}_n, X_1, \dots, X_{m+2})b_j &= (I \otimes Y \otimes X_j \otimes Z)b_j \\ &= (I \otimes Y \otimes X_j \otimes Z)(\sigma_{j-1} \cdots \sigma_n)(\sigma_{n-1} \cdots \sigma_1)(\sigma_1 \cdots \sigma_{n-1})(\sigma_{j-1} \cdots \sigma_n)^{-1} \\ &= (I \otimes X_j \otimes Y^{X_j} \otimes Z)(\sigma_{n-1} \cdots \sigma_1)(\sigma_1 \cdots \sigma_{n-1})(\sigma_{j-1} \cdots \sigma_n)^{-1} \\ &= (X_j \otimes I^{X_j} \otimes Y^{X_j} \otimes Z)(\sigma_1 \cdots \sigma_{n-1})(\sigma_{j-1} \cdots \sigma_n)^{-1} \end{aligned}$$

Since  $I$  contains only 1's, every element of  $I^{X_j}$  is  $1^{m(X_j)} = 1$  and therefore  $I^{X_j} = I$ . Therefore, we get

$$\begin{aligned} (I \otimes Y \otimes X_j \otimes Z)b_j &= (X_j \otimes I \otimes Y^{X_j} \otimes Z)(\sigma_1 \cdots \sigma_{n-1})(\sigma_{j-1} \cdots \sigma_n)^{-1} \\ &= (I \otimes (X_j)^I \otimes Y^{X_j} \otimes Z)(\sigma_{j-1} \cdots \sigma_n)^{-1} \end{aligned}$$

Moreover, since  $I$  is a factorization of 1's,  $m(I)^{-1} = 1$  and therefore we get

$$\begin{aligned} &= (I \otimes X_j \otimes Y^{X_j} \otimes Z)(\sigma_{j-1} \cdots \sigma_n)^{-1} \\ &= (I \otimes Y \otimes X_j \otimes Z) \end{aligned}$$

It remains to be shown that  $(R_1, \dots, R_n, W_1, W_m, H^{-1}, H)b_j$  is equal to

$(R_1^{W_j}, R_n^{W_j}, W_1, W_m, H^{-1}, H)$ . Using the more concise notation  $R = (R_1, \dots, R_n)$ ,  $X = (W_1, \dots, W_{j-1})$ ,  $Y = (W_{j+1}, \dots, W_m, H, H^{-1})$ , and

starting off from the left hand side of the equation, we have:

$$\begin{aligned}
(R_1, \dots, R_n, W_1, W_m, H^{-1}, H)b_j &= \\
&= (R \otimes X \otimes W_j \otimes Y)b_j \\
&= (R \otimes X \otimes W_j \otimes Y)(\sigma_{j-1} \cdots \sigma_n)(\sigma_{n-1} \cdots \sigma_1)(\sigma_1 \cdots \sigma_{n-1})(\sigma_{j-1} \cdots \sigma_n)^{-1} \\
&= (R \otimes W_j \otimes (X)^{W_j} \otimes Y)(\sigma_{n-1} \cdots \sigma_1)(\sigma_1 \cdots \sigma_{n-1})(\sigma_{j-1} \cdots \sigma_n)^{-1} \\
&= (W_j \otimes R^{W_j} \otimes (X)^{W_j} \otimes Y)(\sigma_1 \cdots \sigma_{n-1})(\sigma_{j-1} \cdots \sigma_n)^{-1} \\
&= (R^{W_j} \otimes (W_j)^{R^{W_j}} \otimes (X)^{W_j} \otimes Y)(\sigma_{j-1} \cdots \sigma_n)^{-1}
\end{aligned}$$

Now, we have by hypothesis that  $m(R) = 1$ , and therefore also  $m(R^{W_j}) = 1$ , so we can continue:

$$\begin{aligned}
&= (R^{W_j} \otimes W_j \otimes (X)^{W_j} \otimes Y)(\sigma_{j-1} \cdots \sigma_n)^{-1} \\
&= (R^{W_j} \otimes ((X)^{W_j})^{W_j^{-1}} \otimes W_j \otimes Y) \\
&= (R^{W_j} \otimes (X)^{W_j W_j^{-1}} \otimes W_j \otimes Y) \\
&= (R^{W_j} \otimes X \otimes W_j \otimes Y)
\end{aligned}$$

□

**Lemma 3.4.** *Let  $R$  be the factorization  $(R_1, \dots, R_n)$ ,  $W$  be the factorization  $(W_1, \dots, W_m)$  and  $X$  be the factorization  $(X_1, \dots, X_{m+2})$ . Let  $H$  be an element from  $G_1$ . Then for any  $j$ :  $P_X(R, W, H) \cong_H P_X(R, W, H^{R_j})$*

*Proof.* The proof of this Lemma is very similar to that of the previous lemma. We know from lemma 2.6 that  $P_X(R, W, H) \cong_H P_X(R, W, H^{R_j})$  if and only if there exist a braid  $b \in B_{n+m+2}$  in the stabilizer of  $(\underbrace{1, \dots, 1}_n, X_1, \dots, X_{m+2})$  such that

$$(R_1, \dots, R_n, W_1, \dots, W_m, H^{-1}, H)b = (R_1, \dots, R_n, W_1, \dots, W_m, (H^{-1})^{R_j}, (H)^{R_j})$$

Let us define the braid:

$$c_j = (\sigma_j^{-1} \cdots \sigma_{m+n}^{-1})(\sigma_{m+n+1} \sigma_{m+n+2}^2 \sigma_{m+n+1})(\sigma_{m+n} \cdots \sigma_j), j \leq n$$

and show that it fulfills these conditions. We begin by showing that  $c_j$  is in the stabilizer of  $(\underbrace{1, \dots, 1}_n, X_1, \dots, X_{m+2})$ . Using the shorter notation

$$I = (\underbrace{1, \dots, 1}_j), \quad Z = (\underbrace{1, \dots, 1}_{n-(j+1)}, X_1, \dots, X_m) \quad \text{and} \quad T = (X_{m+1}, X_{m+2}),$$

we see that  $c_j$  is indeed in the stabilizer, because:

$$\begin{aligned} & (\underbrace{1, \dots, 1}_n, X_1, \dots, X_{m+2})c_j \\ &= (I \otimes 1 \otimes Z \otimes T)c_j \\ &= (I \otimes 1 \otimes Z \otimes T)(\sigma_j^{-1} \cdots \sigma_{m+n}^{-1})(\sigma_{m+n+1}^2 \sigma_{m+n+2}^2 \sigma_{m+n+1})(\sigma_{m+n} \cdots \sigma_j) \\ &= (I \otimes (Z)^{1^{-1}} \otimes 1 \otimes T)(\sigma_{m+n+1}^2 \sigma_{m+n+2}^2 \sigma_{m+n+1})(\sigma_{m+n} \cdots \sigma_j) \\ &= (I \otimes Z \otimes 1 \otimes T)(\sigma_{m+n+1}^2 \sigma_{m+n+2}^2 \sigma_{m+n+1})(\sigma_{m+n} \cdots \sigma_j) \\ &= (I \otimes Z \otimes T \otimes 1^T)(\sigma_{m+n+2} \sigma_{m+n+1})(\sigma_{m+n} \cdots \sigma_j) \\ &= (I \otimes Z \otimes T \otimes 1)(\sigma_{m+n+2} \sigma_{m+n+1})(\sigma_{m+n} \cdots \sigma_j) \\ &= (I \otimes Z \otimes 1 \otimes T^1)(\sigma_{m+n} \cdots \sigma_j) \\ &= (I \otimes Z \otimes 1 \otimes T)(\sigma_{m+n} \cdots \sigma_j) \\ &= (I \otimes 1 \otimes Z^1 \otimes T) \\ &= (I \otimes 1 \otimes Z \otimes T) \end{aligned}$$

To conclude the proof, it remains to be proved that

$$(R_1, \dots, R_n, W_1, \dots, W_m, H^{-1}, H)b = (R_1, \dots, R_n, W_1, \dots, W_m, (H^{-1})^{R_j}, (H)^{R_j})$$

Again, using the shorthand notation  $X = (R_1, \dots, R_{j-1})$ ,  $Y = (R_{j+1}, \dots, R_n, W_1, \dots, W_m)$ ,  $H' = (H^{-1}, H)$ , we have:

$$\begin{aligned} & (R_1, \dots, R_n, W_1, \dots, W_m, H^{-1}, H)c_j = \\ &= (X \otimes R_j \otimes Y \otimes H')c_j \\ &= (X \otimes R_j \otimes Y \otimes H')(\sigma_j^{-1} \cdots \sigma_{m+n}^{-1})(\sigma_{m+n+1}^2 \sigma_{m+n+2}^2 \sigma_{m+n+1})(\sigma_{m+n} \cdots \sigma_j) \\ &= (X \otimes Y^{(R_j)^{-1}} \otimes R_j \otimes H')(\sigma_{m+n+1}^2 \sigma_{m+n+2}^2 \sigma_{m+n+1})(\sigma_{m+n} \cdots \sigma_j) \\ &= (X \otimes Y^{(R_j)^{-1}} \otimes H' \otimes R_j^{H'}) (\sigma_{m+n+2} \sigma_{m+n+1})(\sigma_{m+n} \cdots \sigma_j) \end{aligned}$$

We continue by noting that  $m(H') = m((H^{-1}, H)) = 1$ , so we get:

$$\begin{aligned}
& (X \otimes Y^{(R_j)^{-1}} \otimes H' \otimes R_j)(\sigma_{m+n+2}\sigma_{m+n+1})(\sigma_{m+n} \cdots \sigma_j) \\
&= (X \otimes Y^{(R_j)^{-1}} \otimes R_j \otimes (H')^{R_j})(\sigma_{m+n} \cdots \sigma_j) \\
&= (X \otimes R_j \otimes (Y^{(R_j)^{-1}})^{R_j} \otimes (H')^{R_j}) \\
&= (X \otimes R_j \otimes Y^{(R_j)^{-1}R_j} \otimes (H')^{R_j}) \\
&= (X \otimes R_j \otimes Y \otimes (H')^{R_j})
\end{aligned}$$

□

Now that the two Lemmas have been proven, we are ready to tackle Theorem 3.2.

*Proof.* (Theorem 3.2)

We begin by proving that if  $x \in \langle W_1, \dots, W_n \rangle$  then  $P_X(R, W, H) \cong_H P_X(R^x, W, H)$  for any  $H$ . Since  $x \in \langle W_1, \dots, W_n \rangle$ , it can be written as  $x = a_1 \cdots a_\ell$  where  $a_j \in \{W_1, \dots, W_m, W_1^{-1}, \dots, W_m^{-1}\}$ .

We proceed by induction on  $\ell$ . If  $\ell = 0$  then the theorem holds trivially. Next, suppose that the theorem holds for  $\ell$ , and we shall prove that it holds for  $\ell + 1$ . Since  $x$  is the product of  $\ell + 1$  elements from  $\{W_1, \dots, W_m, W_1^{-1}, \dots, W_m^{-1}\}$ , we can write  $x$  as  $x_\ell h_{\ell+1}$ , where  $x_\ell$  is the product of  $\ell$  elements, and  $h_{\ell+1} \in \{W_1, \dots, W_m, W_1^{-1}, \dots, W_m^{-1}\}$ . We know from the induction hypothesis that  $P_X(R, W, H) \cong_H P_X(R^{x_\ell}, W, H)$ . Since  $R_1 \cdots R_n = 1$ , also the product  $(R_1)^{x_\ell} \cdots (R_n)^{x_\ell} = 1$ , and therefore, if  $h_{\ell+1} = W_j$ , we conclude from Lemma 3.3 that  $P_X(R^{x_\ell}, W, H) \cong_H P_X((R^{x_\ell})^{W_j}, W, H) = P_X(R^{x_\ell W_j}, W, H) = P_X(R^x, W, H)$

If on the other hand  $h_{\ell+1} = W_j^{-1}$ , then we know that  $(R_1)^x \cdots (R_n)^x = 1$ , and from Lemma 3.3 we get that  $P_X(R^x, W, H) \cong_H P_X((R^x)^{W_j}, W, H) = P_X(R^{x W_j}, W, H) = P_X(R^{x_\ell W_j^{-1} W_j}, W, H) = P_X(R^{x_\ell}, W, H)$

Now that we have this in hand, we can complete the proof of the theorem. Let  $Y \in N$ . we are going to prove that  $P_X(R, W, H) \cong_H P_X(R, W, H^Y)$ . Since  $Y \in N$  we can write  $Y = (x_1)^{y_1}(x_2)^{y_2} \cdots (x_l)^{y_l}$  where  $x_i \in \{R_1, \dots, R_n, R_1^{-1}, \dots, R_n^{-1}\}$  and  $y_i \in \langle W_1, \dots, W_n \rangle$ .

We prove the theorem by induction on  $l$ . If  $l = 0$ , the theorem holds trivially. Assume that the theorem holds for  $l$  and we shall prove it for

$l + 1$ . Indeed, suppose that  $Y = (x_1)^{y_1}(x_2)^{y_2} \cdots (x_{l+1})^{y_{l+1}}$ .  $Y$  can be factored thus:  $Y = Y_l(x_{l+1})^{y_{l+1}}$ , where  $Y_l = (x_1)^{y_1}(x_2)^{y_2} \cdots (x_l)^{y_l}$ . By the induction hypothesis  $P(R, W, H) \cong_H P(R, W, H^{Y_l})$ .

Suppose now that  $x_{l+1} = (R_j)^{y_{l+1}}$ . We have already proved that  $P_X(R, W, H^{Y_l}) \cong_H P_X(R^{y_{l+1}}, W, H^{Y_l})$ , and from lemma 3.4 we know that  $P_X(R^{y_{l+1}}, W, H^{Y_l}) \cong_H P_X(R^{y_{l+1}}, W, (H^{Y_l})^{R_j^{y_{l+1}}}) = P_X(R^{y_{l+1}}, W, H^{Y_l R_j^{y_{l+1}}}) = P_X(R^{y_{l+1}}, W, H^Y)$ . Since  $y_{l+1} \in \langle W_1, \dots, W_n \rangle$ , by what we have already proved we get that  $P_X(R^{y_{l+1}}, W, H^Y) \cong_H P_X(R, W, H^Y)$ .

On the other hand, if  $x_{l+1} = (R_j^{-1})^{y_{l+1}}$ , since  $y_{l+1} \in \langle W_1, \dots, W_n \rangle$ , by what we initially proved we get that  $P_X(R, W, H^Y) \cong_H P_X(R^{y_{l+1}}, W, H^Y)$  and from lemma 3.4 we get that  $P_X(R^{y_{l+1}}, W, H^Y) \cong_H P_X(R^{y_{l+1}}, W, (H^Y)^{(R_j)^{y_{l+1}}}) = P_X(R^{y_{l+1}}, W, H^{Y R_j^{y_{l+1}}}) = P_X(R^{y_{l+1}}, W, H^{Y_l (R_j^{-1})^{y_{l+1}} R_j^{y_{l+1}}}) = P_X(R^{y_{l+1}}, W, H^{Y_l})$ , and since  $y_{l+1} \in \langle W_1, \dots, W_n \rangle$ ,  $P_X(R^{y_{l+1}}, W, H^{Y_l}) \cong_H P_X(R, W, H^{Y_l})$   $\square$

**Theorem 3.5.** *Let  $G := G_1 \oplus G_2$ , and*

$$\begin{aligned} F_1 &:= ((x_1, 1), \dots, (x_n, 1), (y_1, z_1), \dots, (y_m, z_m)) \\ F_2 &:= ((x'_1, 1), \dots, (x'_n, 1), (y'_1, z_1), \dots, (y'_m, z_m)) \\ K &:= \langle x_1, \dots, x_n, y_1, \dots, y_m \rangle \end{aligned}$$

*Let  $N$  be the normal closure of  $\{x_1, \dots, x_n\}$  inside  $K$ .*

(1) *If the following conditions hold:*

- (a)  $F_1 \cong_H F_2$ ,
- (b)  $z_i \neq 1$ , for all  $i$ ,

*then there exists a permutation  $\pi \in S_n$  such that for any  $i$ ,  $1 \leq i \leq n$ ,  $x'_i$  is conjugated to  $x_{\pi(i)}$  in  $K$  (in particular  $x'_i \in N$ ).*

(2) *If in addition the braid-action stabilizer of  $(z_1, \dots, z_m)$  is contained in the stabilizer of  $(y_1, \dots, y_m)$  then for any  $i$ ,  $1 \leq i \leq m$ , there exists an element  $m_i \in N$  such that  $y'_i = y_i^{m_i}$  (Note that both the stabilizers are subgroups of  $B_m$  and therefore the requirement that one be contained in the other is well defined.)*

*Proof.* (1) Since  $F_1 \cong_H F_2$ , there is a permutation  $\pi_1 \in S_{m+n}$  such that the  $r$ -th element in  $F_2$  is conjugated to the  $\pi_1(r)$ -th element in  $F_1$ . Let  $i$  be a number,  $1 \leq i \leq n$ . We know that  $(x'_i, 1)$  is conjugated in  $G$  to the  $\pi_1(i)$ -th element of the factorization  $F_1$ ,

say  $(a, b)$ . This means that  $x'_i$  is conjugated to  $a$  in  $G_1$  and 1 is conjugated to  $b$  in  $G_2$ , which implies that  $b = 1$ . Now, since  $z_j \neq 1$  for all  $j$ ,  $(x'_i, 1)$  must be conjugated to  $(x_l, 1)$  for some  $l$ ,  $1 \leq l \leq n$ . This in turn implies that  $x'_i$  is conjugated to  $x_l$  in  $K$ , and that  $\pi_1$  induces a permutation on the first  $n$  elements.

- (2) We know from Lemma 2.6 that  $F_1 \cong_H F_2$  if and only if there exists a braid  $b$  in the stabilizer of  $(\underbrace{1, \dots, 1}_n, z_1, \dots, z_m)$  such

that  $(x_1, \dots, x_n, y_1, \dots, y_m)b$  is equal to  $(x'_1, \dots, x'_n, y'_1, \dots, y'_m)$ . From Lemma 2.3 we know that if  $S$  is the stabilizer of  $(z_1, \dots, z_m)$ , then every element in the stabilizer of  $(\underbrace{1, \dots, 1}_n, z_1, \dots, z_m)$  can

be written as  $CBA$  where  $A \in \langle \{\sigma_i | i < n\} \rangle$ ,  $B \in \langle \{A_{jk} | j < n, j \leq k \leq n + m\} \rangle$ ,  $C \in \varphi(S)$ . In particular, we can write  $b = CBA$ , with  $A, B, C$  as above. Since  $C \in \varphi(S)$ , and since we know that the stabilizer of  $(z_1, \dots, z_m)$  is contained in the stabilizer of  $(y_1, \dots, y_m)$ , then  $C$  must be in the stabilizer of  $(x_1, \dots, x_n, y_1, \dots, y_m)$ , so  $(x_1, \dots, x_n, y_1, \dots, y_m)BA = (x'_1, \dots, x'_n, y'_1, \dots, y'_m)$ . In order to prove the theorem, we must prove that for any braid of the form  $BA$ , with  $A \in \langle \{\sigma_i | i < n\} \rangle$  and  $B \in \langle \{A_{jk} | j < n, j \leq k \leq n + m\} \rangle$ , there exists a factorization  $X$  of length  $n$  and  $p_1, \dots, p_m \in N$  such that  $(F_1)BA = X \otimes (y_1^{p_1}, \dots, y_m^{p_m})$ .

We first prove this fact for  $B$ , by induction on the length of  $B$  when written as a product of the generators  $\{A_{jk} | j < n, j \leq k \leq n + m\}$ . If the length is 0, then the theorem is trivial. Assume now that the theorem is true for  $\ell$  and we will prove it for  $\ell + 1$ . Let  $B = B_\ell b_{\ell+1}$ , where  $B_\ell$  is a braid of length  $\ell$  and  $b_{\ell+1}$  is a braid of the form  $A_{ir}^\varepsilon$ , where  $i < n$ ,  $i \leq r \leq n + m$  and  $\varepsilon \in \{\pm 1\}$ . From the induction hypotheses, we conclude that there exists a factorization of length  $n$  and  $p_1, \dots, p_m \in N$ , such that  $(x_1, \dots, x_n, y_1, \dots, y_m)B_\ell = X \otimes (y_1^{p_1}, \dots, y_k^{p_k})$ . From Corollary 1.7 we know that  $A_{ir}^\varepsilon$  acts on every element except the  $i$ -th element of  $X \otimes (y_1^{p_1}, \dots, y_k^{p_k})$  by conjugating it by an element of the normal closure of the  $i$ -th element inside  $K$ . In particular, since  $i < n$ , it acts on all the elements beyond the  $n$ -th one by conjugating them by elements from normal closure of the  $i$ -th element. From the first part of the theorem we know that the  $i$ -th element is in  $N$ , which implies that its normal closure is

also contained in  $N$ , so there is a factorization  $X'$  of length  $n$ , and  $P_1, \dots, P_m \in N$  such that  $X \otimes (y_1^{p_1}, \dots, y_k^{p_k}) A_{ir}^\varepsilon = X' \otimes ((y_1^{p_1})^{P_1}, \dots, (y_k^{p_k})^{P_k}) = X' \otimes (y_1^{p_1 P_1}, \dots, y_k^{p_k P_k})$ . We conclude the induction by noting that for any  $j$ ,  $p_j$  and  $P_j$  are in  $N$ , whence also  $p_j P_j \in N$ .

To complete the proof, we must prove the fact for  $A$ . However, since  $A$  is generated by braids of the form  $\{\sigma_i \mid i < n\}$ , all the elements beyond the  $n$ -th element remain fixed.

□

**Corollary 3.6.** *Let  $R$  be the factorization  $(R_1, \dots, R_n)$ ,  $W$  be the factorization  $(W_1, \dots, W_m)$  and  $X$  be the factorization  $(X_1, \dots, X_{m+2})$ . Let  $H$  and  $H_1$  be elements from  $G_1$ . Assume that the following conditions hold:*

- (1)  $R_1 \cdots R_n = 1$ .
- (2)  $H \in \langle R_1, \dots, R_n, W_1, \dots, W_m \rangle$ .
- (3) For every  $i$ ,  $X_i \neq 1$ .
- (4) The braid-action stabilizer of  $(X_1, \dots, X_{m+2})$  is contained in the stabilizer of  $(W_1, \dots, W_m, H^{-1}, H)$ . (Note that both of the stabilizers are subgroups of  $B_{m+2}$ , therefore the question of whether one is contained in the other is well defined.)

Then the normal subgroup of  $\langle R_1, \dots, R_n, W_1, \dots, W_m \rangle$  generated by  $\{R_1, \dots, R_n\}$  contains a word  $Y$  such that  $H^Y = H_1$  if and only if  $P_X(R, W, H) \cong_H P_X(R, W, H_1)$ .

*Proof.* The *if* part follows trivially from Theorem 3.2. For the *only if* direction, we use Theorem 3.5, which implies that if  $P_X(R, W, H) \cong_H P_X(R, W, H_1)$ , then there is an element  $Y$  in the normal closure of  $\langle R_1, \dots, R_n, W_1, \dots, W_m, H^{-1}, H \rangle$ , generated by  $\{R_1, \dots, R_n\}$  such that  $H^Y = H_1$ . Since  $H \in \langle R_1, \dots, R_n, W_1, \dots, W_m \rangle$ ,  $\langle R_1, \dots, R_n, W_1, \dots, W_m, H^{-1}, H \rangle = \langle R_1, \dots, R_n, W_1, \dots, W_m \rangle$  and  $Y$  is indeed in the normal closure of  $\{R_1, \dots, R_n\}$  in  $\langle R_1, \dots, R_n, W_1, \dots, W_m \rangle$

□



4. THE FUNCTIONS  $FTL_n$ 

We shall now show that the word problem and the Hurwitz equivalence problem are equal in some sense. We first present some well known notation on the free group. A word is said to be *reduced* if there are no two adjacent letters of the form  $x_i x_i^{-1}$  or  $x_i^{-1} x_i$ . A word is *cyclic reduced* if it is reduced and the first and the last letters are not inverses. We say that a word  $A$  has a *root* if there is a word  $w$  and  $n \in \mathbb{Z}, n \neq \pm 1$ , (or equivalently,  $n \geq 2$ ) such that  $A = w^n$ .

We begin by defining the sets:

$$S_1 := \left\{ (F_n, V_m) \left| \begin{array}{l} F_n \text{ is a free group generated by } z_1, \dots, z_n \text{ for some } n, 1 \leq n < \infty \\ V_m = (v_1, \dots, v_m) \in (F_n^*)^m, 0 \leq m < \infty \end{array} \right. \right\}$$

$$NS_1 := \{ (F_n, V_m, a) \mid (F_n, V_m) \in S_1, a \in F_n^* \}$$

Next, we define a function  $FTL_{s1}$  from  $NS_1$  to the set of 1-factorization of the group  $F_2 \oplus F_2$ .

**Definition 4.1** ( $FTL_{s1}$ ).

Let  $(F_n, V_m, a) \in NS_1$ . Let  $F_X$  be the free group with  $n+1$  generators, generated by  $X_1, \dots, X_{n+1}$ , where these symbols are taken to be distinct of the symbols in  $(F_n, V_m, a)$ .

Note that for every  $r_1, r_2 \in \mathbb{N}$   $F_{r_1} \oplus F_{r_2} \subseteq F_2 \oplus F_2$ , so we can chose a family of injections  $i_{(r_1, r_2)} : F_{r_1} \oplus F_{r_2} \rightarrow F_2 \oplus F_2$ . We define  $FTL_{s1}(F_n, V_m, a) := i_{(n, n+1)} P_X(Vf, W, a)$  where  $Vf := (v_1, \dots, v_m, (v_1 \cdots v_m)^{-1})$ ,  $W := (z_1, \dots, z_n, (z_1 \cdots z_n)^{-1})$  and  $X := (X_1, \dots, X_n, (X_1 \cdots X_n)^{-1}, X_{n+1}^{-1}, X_{n+1})$ . We can easily see that  $FTL_{s1}(F_n, V_m, a)$  is a 1-factorization.

**Theorem 4.2.** Let  $(F_n, V_m) \in S_1$ . Let  $a, b$  be words from  $F_n$  and  $N$  be the normal closure of  $V_m$  in  $F_n$ . Then  $FTL_{s1}(F_n, V_m, a) \cong_H FTL_{s1}(F_n, V_m, b)$  if and only if there exists an element  $Y \in N$  such that  $a^Y =_{F_n} b$

*Proof.* It is sufficient to prove that  $P_X(Vf, W, a) \cong_H P_X(Vf, W, b)$  if and only if there exists  $Y$  in the normal closure of  $V_m$  in  $F_n$  such that  $a^Y =_{F_n} b$ . This will follow from Corollary 3.6, once we've confirmed that all the conditions of the corollary hold. But indeed,

- (1) The product of the elements of  $Vf$  is 1

- (2) Every element in  $X$  is different from 1.
- (3) The word  $a$  is in  $F_n$ .
- (4) The stabilizer of  $X$  is contained in the stabilizer of  $W \otimes a^{-1} \otimes a$ , for any  $a$ . This is true because we can construct an homomorphism  $h$  from  $F_X$  to  $G_W$  by  $X_i \mapsto z_i$  for  $1 \leq i \leq n$  and  $X_{n+1} \mapsto a$ . Since  $h$  is a homomorphism, by Lemma 2.2 the stabilizer of  $X$  is contained in stabilizer of  $W \otimes a^{-1} \otimes a = h(X)$

We can therefore conclude from Corollary 3.6 that  $P_X(Vf, W, a) \cong_H P_X(Vf, W, b)$  if and only if there exists an element  $Y$  such that  $a^Y =_{F_n} b$ , where  $Y$  is in the normal closure of  $\{v_1, \dots, v_m, (v_1 \cdots v_m)^{-1}\}$  in  $\langle v_1, \dots, v_m, (v_1 \cdots v_m)^{-1}, z_1, \dots, z_n, (z_1 \cdots z_n)^{-1} \rangle$ , which is equal to the normal closure of  $\{v_1, \dots, v_m\}$  in  $\langle z_1, \dots, z_n \rangle$  (i.e.  $N$ )  $\square$

**Lemma 4.3.**

- (1) If  $a$  and  $b$  are elements of a free group  $F$ , such that  $a^m$  and  $b^n$  commute, then  $a$  and  $b$  must be powers of a common element  $c$ .
- (2) If  $a$  is an element with no root in the free group  $F$ , then its centralizer is  $Z(a) = \{a^k \mid k \in \mathbb{Z}\}$

*Proof.* (1) See [14].

- (2) Obviously  $a$  and  $a^k$  commute for any  $k \in \mathbb{Z}$  so what remains to be shown is that if  $b \in Z(a)$ , then  $b = a^k$  for some  $k \in \mathbb{Z}$ . But indeed, if  $b \in Z(a)$ , then by definition  $a$  and  $b$  commute, which by (1) means that they are both powers of a common element  $c$ , i.e.  $a = c^m$  and  $b = c^n$ . However, since  $a$  has no root, we must conclude that  $m = \pm 1$ , which implies  $c = a^{\mp 1}$ , so in summary we have  $b = c^n = (a^{\mp 1})^n = a^{\mp n}$ , which concludes the proof.  $\square$

**Lemma 4.4.** Let  $F_n$ , ( $n \geq 2$ ) be a free group, and let  $N$  be a non trivial normal subgroup. There exists an element  $A$  in  $N$  such that  $A$  has no root

*Proof.* Let  $y \neq e$  be some element in  $N$ . Since  $N$  is normal we may assume that  $y$  is cyclic-reduced. We shall separate our proof into two cases:

*Case 1:*  $y = (x_i)^n$ .

Without loss of generality we can take  $y = (x_1)^n$ . Now define  $A =$

$x_2^{-1}(x_1)^n x_2(x_1)^n$ . Note that  $A$  is cyclic reduced. We claim that  $A$  has no root. Indeed, assume by negation that there is a cyclic reduce word  $w$  such that for some  $m \geq 2$ ,  $w^m = A$ .  $w$  must contain  $x_2^{-1}$  which means that  $w^m$  must contain  $x_2^{-1}$   $m$  times, which is not the case.

*Case 2:*  $y$  contains at least two different generators.

In this case, we can write our word (perhaps after cyclic permutation) in such a way that it begins and ends with different generators. Without loss of generality we can write the word as  $y = x_1 Z x_2^\varepsilon$  where  $\varepsilon \in \{1, -1\}$ . Denote the length of  $y$  by  $l$ , and define  $A = (x_2^\varepsilon x_1)^{-(l+1)} x_1 Z x_2^\varepsilon (x_2^\varepsilon x_1)^{(l+1)} x_1 Z x_2^\varepsilon$ . Again,  $A$  is cyclic reduced and we claim that it has no root. Suppose, to the contrary, that there is a word  $w$  such that for some  $m \geq 2$   $w^m = A$ . Since our word start with  $(x_2^\varepsilon x_1)^{-(l+1)}$  and also  $x_1$  appears in  $A$ , also  $w$  must start with  $(x_2^\varepsilon x_1)^{-(l+1)}$ . In addition, since our word contain  $(x_2^\varepsilon x_1)^{(l+1)}$  and since  $w$  start with  $x_1^{-1}$ ,  $w$  must contain  $(x_2^\varepsilon x_1)^{(l+1)}$ . Summing up, we see that the length of  $w$  is at least  $4l + 4$ , so the length of  $w^m$  is at least  $m(4l + 4) > 6l + 4$ . However, the length of  $A$  is only  $6l + 4$ , and we have reached a contradiction.  $\square$

Let us define the sets:

$$S_2 := \{(F_n, V_m) \mid (F_n, V_m, a) \in NS, n \geq 2, \exists v \in V_m \text{ such that } v \neq_{F_n} 1\}.$$

$$NS_2 := \{(F_n, V_m, a) \mid (F_n, V_m, a) \in NS_1, (F_n, V_m) \in S_2\}.$$

We shall now construct a function from  $NS_2$  to the set of 1-factorizations in the group  $F_2 \oplus F_2$ , which we denote  $FTL_{s_2}$ .

**Definition 4.5** ( $FTL_{s_2}$ ).

Let  $(F_n, V_m, a) \in NS_2$ , and let  $N$  be the normal closure of  $V_m$  in  $F_n$ .  $V_m$  contains a word which represents a non-trivial element in  $F_n$ , so  $N$  cannot be trivial. Since  $n \geq 2$  and  $N$  is non-trivial, from Lemma 4.4 we conclude that there is a word  $H$  in  $N$  which has no root. We fix such an  $H$  for  $F_n$  and  $V_m$ , and define  $FTL_{s_2}(F_n, V_m, a) = FTL_{s_1}(F_n, V_m, H^a)$

**Theorem 4.6.** Let  $(F_n, V_m) \in S_2$ . Let  $a, b$  be words in  $F_n$  and  $N$  be the normal closure of  $V_m$  in  $F_n$ . Then  $FTL_{s_2}(F_n, V_m, a) \cong FTL_{s_2}(F_n, V_m, b)$  if and only if  $a^{-1}b \in N$ .

*Proof.* By definition  $FTL_{s_2}(F_n, V_m, a) \cong_H FTL_{s_2}(F_n, V_m, b)$  if and only if  $FTL_{s_1}(F_n, V_m, H^a) \cong_H FTL_{s_1}(F_n, V_m, H^b)$ . From Theorem 4.2 we

know that  $FTL_{s1}(F_n, V_m, H^a) \cong_H FTL_{s1}(F_n, V_m, H^b)$  if and only if  $\exists Y \in N$  such that  $(H^a)^Y = H^b$ . Now,  $Y_1 = a^{-1}b$  is readily seen to be a solution of the equation  $(H^a)^Y = H^b$ , so the general solution is  $XY_1$ , where  $X$  commutes with  $H^a$ . However, since  $H$  has no root, neither does  $H^a$ , so from Lemma 4.3 we know that only  $(H^a)^k$  commute with  $H^a$ , and therefore  $Y$  must be equal to  $(H^a)^k a^{-1}b$  for some  $k \in \mathbb{Z}$ . Therefore,  $FTL_{s2}(F_n, V_m, a) \cong FTL_{s2}(F_n, V_m, b)$  if and only if  $(H^a)^k a^{-1}b \in N$  for some  $k \in \mathbb{Z}$ . However,  $H \in N$  and  $N$  is normal in  $F_n$ , so  $(H^a)^k \in N$  for any  $k \in \mathbb{Z}$ . From this we conclude that  $FTL_{s2}(F_n, V_m, a) \cong FTL_{s2}(F_n, V_m, b)$  if and only if  $a^{-1}b \in N$   $\square$

Let us now define the set  $GS$  of all groups with at least two generators and a non trivial relation, and the set

$$GSA := \{(G, a) \mid G \in GS, a \in G^*\}$$

We shall now construct a function from  $GSA$  to the set of 1-factorizations of  $F_2 \oplus F_2$ , which we denote  $FTL_B$ .

**Definition 4.7.**  $FTL_B$

Let  $G := \langle W_1, \dots, W_n \mid R_1, \dots, R_m \rangle$ , and let  $a$  be a word in  $G$ . Let  $F_n$  be the free group with the generators  $z_1, \dots, z_n$ , and let  $\varphi$  be the rewriting function which maps words in  $G$  to words in  $F_n$  by replacing  $W_i$  with  $z_i$  and  $W_i^{-1}$  with  $z_i^{-1}$ . We define  $FTL_B(G, a) = FTL_{s2}(F_n, (\varphi(R_1), \dots, \varphi(R_m)), \varphi(a))$ . This function is well defined, because  $G$  has at least two generators and a non-trivial relation.

**Theorem 4.8.** Let  $G \in GA$  and let  $a, b$  be words in  $G$ . Then  $a =_G b$  if and only if  $FTL_B(G, a) \cong_H FTL_B(G, b)$ .

*Proof.* Let  $N$  be the normal closure of  $\{\varphi(R_1), \dots, \varphi(R_m)\}$  in  $F_n$ . We know that  $a =_G b$  if and only if  $\varphi(a^{-1}b) \in N$ . By Theorem 4.6 we know that this happens if and only if

$$FTL_{s2}(F_n, (\varphi(R_1), \dots, \varphi(R_m)), \varphi(a)) \cong_H FTL_{s2}(F_n, (\varphi(R_1), \dots, \varphi(R_m)), \varphi(a))$$

which, by definition, is equivalent to  $FTL_B(G, a) \cong_H FTL_B(G, b)$   $\square$

**Theorem 4.9.**  $F_2 \oplus F_2 \subseteq B_5 \subseteq B_n, (n > 5)$

*Proof.* [9]  $\square$

**Definition 4.10** ( $FTL_n$ ).

From Theorem 4.9 we know that there exists a family of injections  $ib_n : F_2 \oplus F_2 \rightarrow B_n$ . We define the functions  $FTL_n$  from  $GS$  to the set of 1-factorizations in  $B_n$  by  $FTL_n(G, a) = ib_n(FTL_B(G, a))$

**Corollary 4.11.** *Let  $G \in GS$ , and let  $a, b$  be words in  $G$ . Then  $a =_G b$  if and only if  $FTL_n(G, a) \cong_H FTL_n(G, b)$*

Note that groups with a single generator or with no relations can be changed into the desired form by adding a generator and a relation which equates it to the identity. We can also notice that these groups are the cyclic groups and the free groups, which have uniform algorithms for solving their word problems. Therefore, we may assume that our function is defined on all finitely presented groups.

## 5. THE MAIN RESULTS

The following theorem is our main undecidability result.

**Theorem 5.1.**

- (1) *The Hurwitz equivalence problem for 1-factorizations and  $\Delta^2$ -factorizations in  $B_n$  ( $n \geq 5$ ) is undecidable.*
- (2) *For any  $n \geq 5$  there exist a 1-factorization of  $B_n$ , which we denote by  $u$ , such that the problem of determining whether a 1-factorization  $v$  is Hurwitz-equivalent to  $u$  is undecidable.*
- (3) *For any  $n \geq 5$  there exist a  $\Delta^2$ -factorization of  $B_n$ , denoted by  $u$ , such that the problem of determining whether a  $\Delta^2$ -factorization  $v$  is Hurwitz-equivalent to  $u$  is undecidable.*

*Proof.*

- (1) Follows from (2) and (3)
- (2) Let  $G$  be some finitely presented group with an undecidable word problem. Let  $FTL_n$  be the function into  $\{1\text{-factorizations in } B_n\}$  which we constructed in 4.10. Define  $u := FTL_n(G, 1)$ , a 1-factorization of  $B_n$ , ( $n \geq 5$ ). Suppose now that there exists an algorithm which decides whether a given 1-factorization  $v$  is Hurwitz equivalent to  $u$  or not. In particular, this algorithm would decide whether  $FTL_n(G, 1) \cong_H FTL_n(G, a)$  for

any word  $a$  in  $G$ . By 4.11, this is the same as deciding whether  $a =_G 1$ , which contradicts our choice of  $G$  as having an undecidable word problem.

- (3) Let  $u$  and  $v$  be two 1-factorizations. Since  $\Delta^2$  is central,  $u \cong_H v$  if and only if  $\Delta^2 \otimes u \cong_H \Delta^2 \otimes v$ . However,  $\Delta^2 \otimes u$  and  $\Delta^2 \otimes v$  are  $\Delta^2$ -factorizations, which proves the theorem.

□

**Definition 5.2.** *Let  $u$  be a factorization and  $X$  a set of factorizations.  $u$  is said to be compatible with  $X$  if there exists a finite algorithm which, when given any  $v \in X$ , decides whether  $u \cong_H v$  or not.*

The following theorem gives unrecognizability results.

**Theorem 5.3.**

- (1) *There exists a recursive subset  $X$  of the  $\Delta^2$ -factorizations of  $B_n$   $n \geq 5$ , such that if we take the subset*

$$X_1 = \{u \in X \mid u \text{ is compatible with } X\},$$

*then:*

- (a) *the Hurwitz-equivalence problem on  $X_1$  is undecidable.*
  - (b)  *$X_1$  is not recursively enumerable.*
  - (c) *The problem of determining whether a  $\Delta^2$ -factorization is compatible with  $X$  is unrecognizable.*
- (2) *The problem of determining whether a recursive subset of the  $\Delta^2$ -factorizations of  $B_n$ ,  $n \geq 5$  has a decidable Hurwitz-equivalence problem is unrecognizable.*
- (3) *There exists a recursive subset  $X$  of the  $\Delta^2$ -factorizations such that*
- (a) *the problem of determining whether a recursive subset of  $X$  has solvable Hurwitz equivalence problem is unrecognizable.*
  - (b) *For any  $u$  in  $X$ , the problem of determining whether a recursive subset of  $X$  is compatible with  $u$  is unrecognizable.*

*Proof.* (1) We know that the set of presentations of finitely presented groups is a recursive set. This means that there exists an injective total recursive function  $h$  from the presentations of

finitely presented groups into  $\mathbb{N}$ . We now define the set

$$X := \left\{ \Delta^{2(h(G)+1)} \otimes \Delta^{-2h(G)} \otimes FTL_n(G, A) \mid \begin{array}{l} G \text{ is a finitely presented group} \\ A \text{ is a word in } G \end{array} \right\}$$

Let  $u \in X$ , so  $u$  can be written as  $\Delta^{2(h(G)+1)} \otimes \Delta^{-2h(G)} \otimes FTL_n(G, A)$  where  $G$  is a finitely presented group and  $A$  is a word in  $G$ . Since for any finitely presented group  $G'$  and any word  $A$  from  $G'$  there is no element of the form  $\Delta^{2k}$ ,  $k \in \mathbb{Z}$ , and since  $h$  is injective and  $\Delta^{2k}$  is in the center for any  $k \in \mathbb{Z}$ , the only elements that can be Hurwitz-equivalent to  $u$  are  $\{\Delta^{2(h(G)+1)} \otimes \Delta^{-2h(G)} \otimes FTL_n(G, A) \mid A \text{ is a word in } G\}$ . This set has decidable Hurwitz equivalence problem if and only if  $G$ 's word problem is solvable, so the subset containing all the factorizations for which there exists an algorithm which determines whether they are Hurwitz-equivalent to some factorization from  $X$  is  $\{\Delta^{2(h(G)+1)} \otimes \Delta^{-2h(G)} \otimes FTL_n(G, A) \mid G \text{ is a finitely presented group with solvable word problem and } A \text{ is a word in } G\}$ . We can now prove our theorem.

- (a) Suppose, by contradiction, that there exists an algorithm which solves the Hurwitz-equivalence in this set. This would imply that there is a uniform algorithm to solve all word problems for groups with solvable word problems, which we know is impossible.
  - (b) Assume by negation that  $X_1$  is recursively enumerable. This would mean that there is an algorithm which halts on an element in  $X_1$  if and only if the element is compatible with  $X$ . Now, letting  $G$  be a finitely generated group, we can recognize whether  $G$ 's word problem is solvable by checking whether  $FTL_n(G, 1)$  is compatible with  $X$ . This is again impossible.
  - (c) Suppose that the set of compatible factorizations is recursively enumerated. We know that  $X$  is a recursive set, so their intersection, which is  $X_1$ , is recursively enumerable, contradicting 1b.
- (2) Let  $G$  be a finitely presented group and  $W$  the set of all its words. Determining whether the set  $\{\Delta^2 \otimes FTL_n(G, A) \mid A \in W\}$  has decidable Hurwitz-equivalence problem is equivalent

to determining the problem of whether  $G$ 's word problem is unsolvable, which is known to be an unrecognizable problem.

- (3) We know [8] that there exists a universal finitely presented group  $K$  and a Turing machine  $T$  such that for any finitely presented group  $G$ ,  $T(G)$  is a finite set of words in  $K$  such that  $\langle T(G) \rangle \simeq G$ . We denote  $X := \{FTL_n(K, A) \mid A \text{ is a word in } K\}$ .

- (a) Let  $G$  be a finitely presented group. To determine whether the recursive set

$$\{\Delta^2 \otimes FTL_n(K, A) \mid A \text{ is a word generate by } T(G) \text{ and their inverses}\}$$

has solvable Hurwitz problem is equivalent to determining whether  $G$  has solvable word problem, which is an unrecognizable problem.

- (b) Let  $u \in X$ , so we can write  $u$  as  $\Delta^2 \otimes FTL_n(K, B)$  where  $B$  is a word in  $K$ . Let  $G$  be a finitely presented group. In order to recognize whether  $G$  has solvable word problem we can ask whether for any word  $C \in \langle T(G) \rangle$  we can tell whether  $C =_K 1$ , which is equivalent to asking whether  $CB =_K B$ . So in conclusion, if we consider the recursive set which is also subset of  $X$ :

$$\{\Delta^2 \otimes FTL_n(K, AB) \mid A \text{ is a word generated by } T(G) \text{ and their inverses}\},$$

recognizing whether this set is compatible with  $u = \Delta^2 \otimes FTL_n(K, B)$  is equivalent to recognizing whether  $G$  has solvable word problem, which is an unrecognizable problem.

□

**Remark 5.4.** *All the theorems in this chapter will hold if we take instead  $B_n$  any group including  $F_2 \oplus F_2$  and instead  $\Delta^2$  an element from the center. We give now a proof for one theorem but the rest of the proofs are similar*

**Theorem 5.5.** *let  $G$  be a group including  $F_2 \oplus F_2$  and  $g \in Z(G)$  then the Hurwitz equivalence problem for  $g$  – factorization is undecidable*

*Proof.* since  $F_2 \oplus F_2 \subseteq G$  there is a injective homomorphism  $i : F_2 \oplus F_2 \rightarrow G$  we shell define a function as in the braid group from  $GSA$  to the  $g$  – factorization by  $g \otimes i(FTL_{s2}(K, a)$  where  $K$  is a finitely presented group and a word from this group. We could see that the



hurwitz equivalence problem for  $g$  – *factorization* is undecidable while the proof is very similar from now on to the case of the braid group.  $\square$

## REFERENCES

- [1] S. I. Adian, Finitely presented groups and algorithms, Dokl. Akad. Nauk SSSR 117, 9-12 (1957).
- [2] S. I. Adian, The unsolvability of certain algorithmic problems in the theory of groups, Trudy Moskov. Mat. Obsc. 6, 231-298 (1957).
- [3] E. Artin, The theory of braids. American Scientist 38, (1950). 112–119
- [4] J. Birman, Braids, links and mapping class groups (Princeton University Press, 1975) .
- [5] W. W. Boone, Certain simple unsolvable problems in group theory, I, II, III, IV, V, VI, Nederl. Akad. Wetensch. Proc. Ser. A57, 231-237, 492- 497 (1954), 58, 252-256, 571-577 (1955), 60, 22-27, 227-232 (1957).
- [6] W. W. Boone and H. Rogers Jr., On a problem of J.H.C. Whitehead and a problem of Alonzo Church, Math. Scand. 19, 185-192 (1966).
- [7] W. Buaru *Über zopf-invarianten* .Abh .Math. Sem. Univ. Hamburg, 9, 117 – 124 (1932)
- [8] G. Higman, Subgroups of finitely presented groups, Proc. Royal Soc. London Ser. A 262, 455-475 (1961).
- [9] T. A. Makanina, The occurrence problem for the braid group  $B_n$  with  $n > 5$ , Mat. Zametki 29 (1981), 31-33; English transl., Math. Notes 29 (1981), 16-17.
- [10] C. F. Miller III, (5-MELB) Decision problems for groups—survey and reflections. Algorithms and classification in combinatorial group theory (Berkeley, CA, 1989), 1–59, Math. Sci. Res. Inst. Publ., 23, Springer, New York, 1992.
- [11] P. S. Novikov On the algorithmic unsolvability of the word problem in group theory, Trudy Mat. Inst. Steklov 44, 1-143 (1955).
- [12] S. Vik. Kulikov ; M. Teicher, Braid monodromy factorizations and diffeomorphism types. (Russian) Izv. Ross. Akad. Nauk Ser. Mat. 64 (2000), no. 2, 89–120; translation in Izv. Math. 64 (2000), no. 2, 311–341
- [13] M. O. Rabin, Recursive unsolvability of group theoretic problems, Annals of Math. 67,
- [14] Lyndon, C. Roger ; Paul E. Schupp; Combinatorial group theory. Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 89. Springer-Verlag, Berlin-New York, 1977

DEPARTMENT OF MATHEMATICS AND STATISTICS, BAR-ILAN UNIVERSITY,  
RAMAT-GAN 52900, ISRAEL

*E-mail address:* Liberman@macs.biu.ac.il

DEPARTMENT OF MATHEMATICS AND STATISTICS, BAR-ILAN UNIVERSITY,  
RAMAT-GAN 52900, ISRAEL

*E-mail address:* teicher@macs.biu.ac.il